

Humboldt Mill Aquatic Survey Report 2014

December 1, 2014

Prepared for:

Eagle Mine

Prepared by:

**ADVANCED ECOLOGICAL MANAGEMENT
22071 7 Mile Road
Reed City, MI 49677**

AeM

**ADVANCED
ECOLOGICAL
MANAGEMENT**

Table of Contents

1.0 EXECUTIVE SUMMARY	1
2.0 INTRODUCTION	2
3.0 STUDY SITE	2
4.0 METHODS	6
4.1 Fish Collection	6
4.2 Macroinvertebrates	9
4.3 Stream Habitat Evaluation	9
5.0 RESULTS AND DISCUSSION	11
5.1 Streams	11
5.1.1 Station 1	11
5.1.2 Station 5	12
5.1.3 Station MBER1	12
5.1.4 Station MBER2	13
5.2 Lake Lory and Wetland Complex EE	14
5.2.1 Lake Lory	14
5.2.2 Wetland Complex EE	16
REPORT FIGURES	19
Figure 1-1. Location of Humboldt Mill, Marquette County, Michigan	20
Figure 1-2. Stream Sample and Wetland Complex EE Locations	21
Figure 1-3. Lake Lory Aquatic Survey Locations	22
REPORT TABLES	23
Table 1-1. Summary of the Procedure 51 Macroinvertebrate and Aquatic Habitat Scores for all Stream Stations, 2014	24
Table 3-1. Stream Station Location Description	25
Table 5-2. Stream Macroinvertebrate Collection Data – Stations 1, 5, MBER1 and MBER2	27
Table 5-2 (Continued). Stream Macroinvertebrate Collection Data – Stations 1, 5, MBER1 and MBER2	28
Table 5-3. Stream Macroinvertebrate Scores and Community Ratings – Stations 1, 5, MBER1 and MBER2	29
Table 5-4. Stream Habitat Scores and Ratings – Stations 1, 5, MBER1 and MBER2	30
Table 5-5. Stream Station Dimensions	31
Table 5-6. Average Water Quality Parameters – Stations 1, 5, MBER1, MBER2 and Surface Water Bodies	32
Table 5-7. Lake Lory Fish Collection Data	33
Table 5-8. Lake Lory Fish Size	34
Table 5-9. Lake Lory Macroinvertebrates	35
Table 5-10. Wetland Complex EE Macroinvertebrates	36
PHOTOGRAPHS	37
Photograph C-1. Station 1 – Upstream Extent. View looking south, downstream	38
Photograph C-2. Station 1 – Downstream Extent. View looking west, upstream	38
Photograph C-3. Station 5 – Upstream Extent. View to north, downstream	39
Photograph C-4. Station 5 – Downstream Extent. View to south, upstream	39
Photograph C-5. Station MBER1 – Upstream Extent. View to east, downstream	40
Photograph C-6. Station MBER1 – Downstream Extent. View to west, upstream	40
Photograph C-7. Station MBER2 – Upstream Extent. View to south, downstream	41
Photograph C-8. Station MBER2 – Downstream Extent. View to north, upstream	41
Photograph C-9. Lake Lory. View to north	42

Photograph C-10. Lake Lory. View to south.....	42
Photograph C-11. Wetland Complex EE North of HTDF. View to northwest.....	43
Photograph C-12. Wetland Complex EE North of HTDF. View to north.....	43

List of Abbreviations, Acronyms, and Symbols

AEM	Advanced Ecological Management LLC
CPUE	Catch-per-unit-effort
°C	Degrees Celcius
<i>t</i>	Duration of time
ft	Feet
gpm	Gallons per minute
GLEAS	Great Lakes and Environmental Assessment Section
HTDF	Humboldt Tailings Disposal Facility
KEMC	Kennecott Eagle Minerals Company
MDEQ	Michigan Department of Environmental Quality
µS/cm	Microsiemens per centimeter
mg O ₂ /L	Milligrams of oxygen per liter of water
ml	Milliliters
max	Maximum
min	Minimum
N	North
n.m.	Not measured
<i>N</i>	Number of fish caught
pH	Measure of acidity or alkalinity of a solution
P-51	Procedure Number 51
R	Range
<i>n</i>	Sample size
Sec	Section
<i>s</i>	Standard deviation
T	Township
W	West

1.0 EXECUTIVE SUMMARY

Advanced Ecological Management, LLC (AEM) conducted an aquatic survey in June 2014 at the Humboldt Mill site for Lundin Mining Corporation's Eagle Mine. The Humboldt Mill site is located in the Upper Peninsula of Michigan, Marquette County as shown on Figure 1-1. An aquatics survey at four stream stations included fish, macroinvertebrate, and habitat community ratings according to the metrics outlined in the Great Lakes and Environmental Assessment Section (GLEAS) Procedure Number 51 (P-51), a survey protocol for wadable streams and rivers. Additional aquatics surveys at Lake Lory, and a wetland complex located northeast of the Humboldt Tailings Disposal Facility were also conducted. Sampling survey locations are shown on Figures 1-2 and 1-3.

A summary of the fish, macroinvertebrate, and habitat ratings for the five stream stations are displayed in the table below. All five stations were rated as “poor” fish communities and “acceptable” macroinvertebrate communities. Stream habitat was considered “excellent” in all stations except Station 5, where habitat was rated as “good”.

	Station 1	Station 5	Station MBER1	Station MBER2
Fish Community	Poor	Poor	Poor	Poor
Macroinvertebrate Community	Acceptable	Acceptable	Acceptable	Acceptable
Stream Habitat	Excellent	Good	Excellent	Excellent

No threatened or endangered species of fish or macroinvertebrates were observed within the five stream sample stations (Michigan Natural Features Inventory, 2014).

The fish community in Lake Lory was predominately comprised of warm water species such as perch, bluegill, largemouth bass, and white suckers, and the fish community in Wetland Complex EE was comprised of fathead minnows. No threatened or endangered fish species were observed in Lake Lory or Wetland Complex EE (Michigan Natural Features Inventory, 2014).

2.0 INTRODUCTION

Humboldt Mill is used to process ore that is hauled in from the nearby Eagle Mine. Eagle Mine is conducting annual aquatic surveys of waters in the vicinity of the Humboldt Mill as part of meeting requirements R 425.202 (2) (y) of Michigan's Natural Resources and Environmental Protection Act 451 of 1994 as amended Part 632. This aquatics survey was conducted by AEM in June 2014 to provide a description of the aquatic communities in the waters surrounding the Humboldt Mill. Previous aquatic surveys of the area were conducted by AEM in 2006, 2007, and 2008 (AEM, April 2007a; AEM October 2007b; AEM October 2007c; AEM, April 2008; AEM, March 2009). The objectives of this survey were to provide a general characterization of aquatic communities and record any threatened or endangered aquatic species encountered. This site-specific survey as well as background information, provides data intended to satisfy mine permit requirements (Permit Condition J-14). This report is based on evaluations of fish, macroinvertebrates, and aquatic habitat that were surveyed during June 2014.

3.0 STUDY SITE

The Humboldt Mill property is a former iron-ore mine and ore processing facility located southeast of Champion, Michigan. The mill property and study area are located in Sections 1, 2, 10, 11, 12, 13, and 14 of Humboldt Township (Township 47 North, Range 29 West), in the vicinity of Highway 41 and east of County Road 95, Marquette County, Michigan (Figure 1-1).

The aquatic investigations conducted by AEM included surveys of Lake Lory, a pond located in the southern extent of a wetland complex located just northeast of the Humboldt Tailings Disposal Facility (herein referred to as Wetland Complex EE), two locations on the Middle Branch Escanaba River, a tributary to the Middle Branch Escanaba River, and one stream in the vicinity of the mill, which is a tributary to the Black River. Stream sample stations and other water body survey locations are illustrated in Figures 1-2 and 1-3.

Altogether, the aquatics sample stations included four stream locations, one lake, and one wetland. These sample stations were chosen based on their proximity to the

processing facility, their ability to provide representative habitat and biological diversity information, and their potential to be impacted by the project.

The following is a description of individual sampling stations for the aquatics portion of the survey:

Station 1: Station 1 is located on the West Branch of the Black River that flows south from a former iron-ore tailings basin (Figure 1-2). The upper end of Station 1 is located approximately 30 feet downstream from an unnamed east-west road that connects to County Road 601, and the stations continue downstream (south) 100 feet. The station location was selected downstream of the road crossing to minimize the influence of beaver impoundments that are located upstream of the road crossing. Station 1 is a low-gradient reach of the Black River.

The substrate was predominantly comprised of organic matter, silt, and woody debris. The predominant riparian vegetation consisted of tussock sedge (*Carex stricta*). Woody shrubs such as sweet gale (*Myrica gale*), speckled alder (*Alnus rugosa*), and sandbar willow (*Salix exigua*) were also observed along the stream bank within the station (Photographs C-1 and C-2).

Station 5: Station 5 is located along the right-of-way of Wolf Road, (Marquette County Road FX) approximately 1,200 feet south of the Middle Branch Escanaba River. The upstream extent of the station is located approximately 20 feet downstream of an abandoned railroad grade. Station 5 is approximately 180 feet in length and flows northeast to the downstream extent, approximately 15 ft upstream of Wolf Road (Figure 1-2). Station 5 drains an adjacent wetland to the west, which is connected to Wetland EE and nearby HTDF.

The substrate of the stream was predominantly comprised of sand, silt, and organic matter. The streambanks were well vegetated with speckled alder overhanging the stream channel, and the predominant herbaceous component was reed canarygrass (*Phalaris arundinacea*; Photographs C-3 and C-4).

Station MBER1: Station MBER1 is located on the Middle Branch Escanaba River northeast of the Humboldt Mill. The downstream extent of the station is located approximately 2,680 feet upstream (north) of Highway 41 (Figure 1-2). Station MBER1 is approximately 1,000 feet long and the stream flows east at this location.

The substrate was predominantly comprised of gravel, silt, sand, and organic matter. The predominant riparian vegetation within Station MBER1 consisted of tussock sedge and speckled alder (Photographs C-5 and C-6). Submerged pondweed (*Potamogeton* sp.) was abundant throughout the stream channel.

An abandoned railroad grade comprised a portion of the south streambank along the approximately 500 feet of the station. The streambank along the railroad grade was vegetated and one culvert connects a wetland located south of the river channel to the river near the upstream extent of the station.

Station MBER2: Station MBER2 is located on the Middle Branch Escanaba River approximately 3.1 miles northeast of the Humboldt Mill and upstream of any potential water that may discharge from the surface of the Humboldt Mill area (Figure 1-2). The downstream extent of the station is located immediately upstream of the former Lake Superior and Ishpeming Railroad bridge crossing that now serves as a rails-to-trails bridge for snowmobiles and off road vehicles. Station MBER2 is approximately 1,000 feet long and the stream flows south at this location.

The substrate was predominantly comprised of gravel, sand, silt, and organic matter. The predominant riparian vegetation within Station MBER2 consisted of speckled alder and tussock sedge (Photographs C-7 and C-8).

Lake Lory: Lake Lory is approximately 128 acres in size and is located approximately 1.3 miles south of Highway 41 and approximately 1.9 miles east of County Road 95 (Figures 1-2 and 1-3). Lake Lory drains into the East Branch of the Black River (Figure 1-3) along its southwestern boundary where an earthen berm controls the hydrology of the lake. Ponds located along the northern boundary of the lake drain into the lake and

historically functioned as a tailings basin for the iron-ore mine. Photographs C-9 and C-10 display views of the lake.

Wetland Complex EE: The pond located within Wetland Complex EE is located approximately 167 feet northeast of HTDF (Figure 1-2). The hydrology of the wetland complex is influenced by the presence of Highway 41 and an abandoned railroad grade, both of which bisect the wetland from east to west and direct water movement through numerous culverts. Water generally moves through the wetland complex in a northerly direction from the vicinity of the HTDF towards Highway 41 and to the Middle Branch Escanaba River. Vegetation in the wetland complex is predominantly comprised of emergent and scrub-shrub wetland vegetation, including cattails (*Typha* sp.) and speckled alder (Photographs C-11 and C-12).

Table 3-1 provides a summary of the location information for the four stream sample stations identified above. Depiction of the sampling locations for the Lake Lory and Wetland Complex EE appear on Figures 1-2 and 1-3.

4.0 METHODS

To characterize the quality of the streams within the vicinity of the mill, flowing and wadable water bodies were sampled according to the Michigan Department of Environmental Quality's (MDEQ) Surface Water Quality Division *Procedure #51 Survey Protocols for Wadable Rivers* (P-51; MDEQ, 2008). The P-51 is a rapid bioassessment protocol that is used to evaluate stream quality based on fish, macroinvertebrates, and stream habitat characteristics.

Surface waters were sampled to characterize the fish communities and to provide a general description of the surface-water aquatic habitat. Lake Lory and Wetland Complex EE (Figures 1-2 and 1-3) were sampled using a variety of aquatic sampling methods, including the use of electroshocking gear, nets, and a sediment grabbing device to collect macroinvertebrates from unwadable water bodies.

4.1 Fish Collection

A backpack electroshocker was used in narrow (approximately ≤ 15 feet) or difficult-to-access stations (e.g., areas with abundant woody debris). A barge-mounted electroshocker was used to sample Stations MBER1 and MBER2, which were deep and wide enough to permit the passage of the barge unit. On Stations MBER1 and MBER2, one pass was conducted in an upstream direction and the duration of electroshocking charge time was recorded for each pass. On Stations 1 and 5, a multi-pass removal was conducted (Van Deventer and Platts, 1983). For each electrofishing pass, stunned fish were placed in a live well for identification and enumeration. Following collection and fish identification of all fish in each station survey, enumerated fish were released within each station.

Stream fish data were analyzed according to P-51 metrics to produce a "fish score" that was used to rate the fish community as poor, neutral (acceptable), or excellent quality. There are ten metrics used to evaluate the fish community diversity and they are listed in the P-51 methodology (MDEQ, 2008). The score for each metric can range from -1 to $+1$ and are described as follows:

- -1 indicates the community is performing outside of (minus) two standard deviations from the average conditions found at excellent sites;
- 0 indicates the community is performing between the average condition and (minus) two standard deviations from the average condition found at excellent sites, and;
- +1 indicates the community is performing better than the average condition found at excellent sites.

The summation of the fish scores can range from -10 to +10 using these metrics. Stations that score from +5 to +10 are considered excellent. Stations that score from -5 to -10 or have less than 50 fish are considered poor, while stations that score from -5 to +5 (including zero) are considered acceptable in fish community structure (MDEQ, 2008).

Lake, pond, and wetland sites were not evaluated for fish using the P-51 procedure because the approach is designed for wadable streams (MDEQ, 2008). Since fish collection gear is not equally selective among sizes and species of fish, a combination of sampling gear, including a boom shocker, a backpack shocker, experimental gill nets, and fyke nets were used to capture fish within Lake Lory and Wetland Complex EE. Gear and sample locations within Lake Lory were selected to provide a broad sampling coverage of aquatic habitat and to minimize the bias created by gear selectivity and avoidance by fish.

A boat-mounted shocking unit and generator (boom shocker) were used to collect fish in shallow water, near-shore areas of Lake Lory (Figure 1-3). Pulsed direct current was used during the survey to minimize trauma to the fish. Electroshocking duration was automatically recorded as the total seconds of electricity that was discharged from the boom shocker in each transect. Boom shocking was conducted at night, which is more effective than shocking during daylight hours (Smith-Root, 2004).

Multiple panel monofilament gill nets of varying mesh size were fished in several locations throughout Lake Lory (Figure 1-3). Each gill net consisted of five, 6- by 25-foot panels ranging from 1.5- to 6-inch stretch mesh. The gill nets were set overnight and were fished for approximately 12 hours.

Fyke nets were constructed of 0.125-inch "Ace"-type nylon mesh coated with green latex net dip. The lead was 15 feet-long and 2-feet high. The frame and the cab were 10-feet long when fully extended. The frame section is formed by two rectangular spring-steel frames that are 2-feet high 4-feet wide. The cab is constructed of two 2-foot diameter steel hoops. The fyke nets were placed throughout Lake Lory and were fished overnight for approximately 24 hours (Figure 1-3).

Wetland Complex EE was sampled in a wadable pond located approximately 450 feet south of Highway 41, and approximately 1,500 feet west of the intersection between Highway 41 and Wolf Road (Figure 1-2). Sampling was conducted with a backpack electroshocker along the south and west shoreline on the edges of dense vegetation.

Catch-per-unit-effort (CPUE) is used as an index of fish abundance. Fish sampling efforts were standardized to units consistent with the Michigan Department of Natural Resources sampling protocol (Schneider et al., 2000). CPUE for the following gear was estimated as follows:

$$CPUE = \frac{N}{t}$$

Where,

N = number of fish caught

t = sample duration in hours (boom shocker)

t = sample duration in net nights (experimental gill net)

t = sample duration in net nights (hoop net).

As part of the enumeration process, the species, length, weight, and number of fish captured were recorded. One representative of each species that was not identifiable in the field was placed in a voucher jar containing 10% formalin for later identification. Each voucher jar was labeled according to the sample location and date. Fish were identified to species using various taxonomic references (Bailey et al., 2003; Coon, 2001; Becker, 1983). The Marquette County Element List (Michigan Natural Features

Inventory, 2014) was reviewed to determine if any threatened, endangered, or special concern aquatic species occurred within the vicinity of the project.

4.2 Macroinvertebrates

Macroinvertebrate sampling was conducted within all stream stations. Upon completion of fish sampling, aquatic macroinvertebrates, including mussels and decapods (crayfish), were collected within each station using D-framed kick-nets (Merritt et al., 1996). Stations were sampled for 45 minutes using two kick-nets (total sample time = 1.5 hours) and samples were collected in all habitat types within each station to characterize the macroinvertebrate community. Collected specimens were stored in 500 ml plastic wide-mouth jars containing 70% ethanol, and were identified using various taxonomic references (Bright, 2014; Merritt et al., 2008; Cummings and Mayer, 1992; Pennak, 1990).

The wadable stream macroinvertebrate data were analyzed according to nine metrics identified in the P-51 methodology. The sum of the macroinvertebrate scores can range from -9 to +9; and like the fish community, are graded as excellent, acceptable (slightly impaired), or poor according to the summation of the metric scores.

Aquatic macroinvertebrate sampling was conducted along the shoreline of Lake Lory and Wetland Complex EE using a D-frame kick-net, and a PONAR sediment-grabbing device was used in Lake Lory to collect macroinvertebrates within the water basin (Figure 1-3).

4.3 Stream Habitat Evaluation

Riparian and instream habitats were qualitatively described for each station. A description of stream morphology included run/riffle/pool/shallow pool configurations, substrate, substrate embeddedness, instream cover, vegetation, flow stability, and bank stability. Stream habitat was rated as excellent, good, marginal, or poor based on P-51 scores interpreted from 10 habitat metrics. Habitat was rated according to the following P-51 habitat scores (MDEQ, 2008):

Habitat Characterization	Total Point Score
1. Excellent	> 154
2. Good	105 – 154
3. Marginal	56 – 104
4. Poor	< 56

Wetted stream width was measured at the downstream extent, upstream extent, and middle of each sample station to describe the physical dimensions of each stream sample station at the time of the survey. The average depth of the sample station was determined from stream depth measured at 20 percent of the channel width, 80 percent of the channel width, and 50 percent of the channel width of the downstream extent, upstream extent, and middle of the station. Stream flow was measured using a Marsh McBirney FLO-MATE 2000™.

Photographs were taken at each station to illustrate the conditions during the sampling period. Water temperature, pH, conductivity, and dissolved oxygen were measured as part of the stream habitat evaluation. These water quality parameters were measured using a Yellow Springs Instrument Professional Plus water quality meter.

5.0 RESULTS AND DISCUSSION

The aquatic sampling of the Humboldt Mill vicinity was conducted from June 6, 2014 through June 10, 2014. Station 1 was sampled on June 10, 2014, Station 5 was sampled on June 9, 2014, and Stations MBER1 and MBER2 were sampled on June 6, 2014. Sampling of Lake Lory and Wetland Complex EE were conducted from June 7, 2014 through June 9, 2014.

5.1 Streams

All streams sampled within the project vicinity are located in a region that is classified as a Northern Lakes and Forests ecoregion (MDEQ, 2008). Therefore, all P-51 scoring was based on metrics that relied on typical data from this ecoregion.

5.1.1 Station 1

A total of 31 fish were observed in Station 1. The dominant species included northern redbelly dace (*Phoxinus eos*), Central mudminnows (*Umbra limi*) and pearl dace (*Margariscus margarita*; Table 5-1). The CPUE estimate was 1.4 fish/minute of backpack electroshocking for the reach. Because fewer than 50 fish were collected from Station 1, the fish community was rated as “poor”. No threatened, endangered, or special concern fish species were observed in Station 1.

A total of 76 macroinvertebrates were collected from Station 1, where Amphipoda (scuds), Dipterans (true flies) and Trichopterans (caddisflies) were the most abundant organisms (Table 5-2). The macroinvertebrate community was rated as “acceptable” (Table 5-3), and no threatened, endangered, or special macroinvertebrates were observed.

Station 1 was classified as a glide/pool habitat based on use of the P-51 protocol. The habitat was rated as “excellent” based on the total habitat metric score according to the P-51 protocol (Table 5-4).

The average width of Station 1 was 7.1 feet (standard deviation; $s = 3.0$ feet; sample size; $n = 3$) and the average depth was 1.4 feet ($s = 0.5$ feet; $n = 9$; Table 5-5). The discharge within Station 1 was 861 gallons per minute on June 10, 2014. Water

temperature within Station 1 was the coldest observed temperature (13.6°C) and lowest dissolved oxygen (5.0 mg O₂/L) among the Humboldt sample stations (Table 5-6).

5.1.2 Station 5

Station 5 was sampled using a backpack electroshocker where only one Central mudminnow was collected, with an observed CPUE of 0.1 fish/minute of backpack electroshocking. Because fewer than 50 fish and no salmonids were collected from Station 5, the fish community was rated as “poor”. No threatened, endangered, or special concern fish species were observed in Station 5.

A total of 392 macroinvertebrates were collected from Station 2, where Isopoda (sowbugs) were the most abundant organism followed by flies and caddisflies (Table 5-2). The macroinvertebrate community was rated as “acceptable” (Table 5-3) and no threatened, endangered, or special concern macroinvertebrate species were observed.

Station 5 was evaluated as a glide/pool habitat and was rated as “good” based on the P-51 total habitat metric score (Table 5-4). The average width of Station 5 was 3.5 feet ($s = 1.6$ feet; $n = 3$; Table 5-5) and average depth was 0.4 feet ($s = 0.2$ feet; $n = 9$; Table 5-5). The discharge within Station 5 was 95 gallons per minute on June 09, 2014 (Table 5-6). Conductivity (230 μ S/cm) was the highest among stream sample stations (Table 5-6). Although the presence of hydrocarbons are not typically surveyed as part of a P-51, an odor of gasoline, or some other hydrocarbon-based product was noted while conducting the survey of Station 5 when sediments were disturbed from the fish and macroinvertebrate portions of the survey.

5.1.3 Station MBER1

A total of 17 fish representing 7 taxa were observed with an estimated CPUE of 0.3 fish/minute of tote-barge electroshocking in Station MBER1. Northern pike (*Esox lucius*) and white suckers (*Catostomus commersonii*) were the most abundant fish within this station (Table 5-1).

The Middle Branch Escanaba River is classified as a designated trout stream by the State of Michigan (Michigan Department of Natural Resources Fisheries Order 210.04,

2014). Because fewer than 50 fish were collected and no salmonids were collected from Station MBER1, the fish community was rated as “poor”. No threatened, endangered, or special concern fish species were observed in Station MBER1.

A total of 250 macroinvertebrates were collected from Station MBER1, where caddisflies, were most abundant followed by Ephemeropterans (mayflies) and sowbugs (Table 5-2). The total macroinvertebrate community was rated as “acceptable” (Table 5-3), and no threatened, endangered, or special concern macroinvertebrates were observed.

Station MBER1 was classified as a glide/pool habitat and was rated as “excellent” according to the P-51 total habitat metric score (Table 5-4). The average width of Station MBER1 was 36.9 feet ($s = 2.6$; $n = 3$) and average depth was 3.2 feet ($s = 0.7$ feet; $n = 9$; Table 5-5).

The discharge within Station MBER1 was 29,096 gallons per minute on June 6, 2014. Conductivity (73 $\mu\text{S}/\text{cm}$), and pH (6.8) were among the lowest values recorded among the Humboldt sample stations as part of this aquatic survey (Table 5-6).

5.1.4 Station MBER2

Station MBER2 was also sampled using a barge-mounted electroshocker where a total of 19 fish representing 9 taxa were observed. Mottled sculpin (*Cottus bairdii*) were the most frequently observed fish, followed by common shiners (*Notropis cornutus*) and central mudminnows (Table 5-1). The CPUE estimate for Station MBER2 was 0.3 fish/minute of tote-barge electroshocking.

Because fewer than 50 fish and no salmonids were collected from Station MBER2, the fish community was rated as “poor”. No threatened, endangered, or special concern fish species were observed in Station MBER2.

A total of 401 macroinvertebrates were collected from Station MBER2. Dipterans were the most abundant organisms followed by caddisflies and mayflies (Table 5-2). The

macroinvertebrate community was rated as “acceptable” (Table 5-3), and no threatened, endangered, or special concern macroinvertebrate species were observed.

Station MBER2 was evaluated as a glide/pool habitat and was rated as “excellent” based on the P-51 total habitat metric score (Table 5-4). The average width of Station MBER2 was 23.3 feet ($s = 0.4$ feet; $n = 3$) and average depth was 1.6 feet ($s = 0.7$ feet; $n = 9$; Table 5-5). Discharge within Station MBER2 was 13,828 gallons per minute on June 6, 2014 (Table 5-6). Water temperature was the warmest (20.3°C) and conductivity was the lowest (36 $\mu\text{S}/\text{cm}$) among all stream sample stations (Table 5-6).

5.2 Lake Lory and Wetland Complex EE

5.2.1 Lake Lory

Six fyke nets were fished in nearshore areas throughout Lake Lory for approximately 24 hours for each net (Figure 1-3). Sampling took place from June 7, 2014 through June 9, 2014. Fyke nets were typically set mid-day and were checked at the same time the following day for fish.

Gill nets were fished in three locations throughout the lake for approximately 12 hours of soak time for each net (Figure 1-3). Gill nets were fished from June 7 through June 9, 2014.

Boom shocking was conducted along five transects that were situated parallel to the shoreline (Figure 1-3). Transect lengths ranged from a minimum of 257 feet to a maximum of 483 feet.

A total of 227 fish were collected from Lake Lory including 11 different taxa (Table 5-7). Largemouth bass (*Micropterus salmoides*) were the most frequently observed species, followed by bluegills (*Lepomis macrochirus*) and yellow perch (*Perca flavescens*) among all sample gear. No threatened, endangered, or special concern fish species were observed in Lake Lory.

Most of the fish were collected using a boom shocker (Table 5-7). The average total CPUE for the boom shocker was 138 fish/hour and varied among transects.

The total CPUE for fyke nets varied by net location, and ranged from a minimum of 3 fish per net night in fyke nets 1 and 5, to a maximum of 17 fish per net night in fyke net 2. The fyke nets were the most effective means for capturing small minnow species, such as blacknose shiners (*Notropis heterolepis*) and brassy minnows (*Hybognathus hankinsonii*). The total number of fish captured in gill nets was 21 (Table 5-7), and ranged from zero fish in gill nets 3 to 14 fish in gill net 2.

The average length of largemouth bass was 12.0 inches ($s = 1.9$ inches; $n = 79$) and the average weight was 0.8 pounds ($s = 0.4$ pounds; $n = 79$; Table 5-8). Bluegill ranged in length from 0.4 to 9.3 inches, with an average length of 5.5 inches ($s = 2.0$ inches; $n = 69$), and an average weight of 0.1 pounds ($s = 0.1$ pounds; $n = 69$ Table 5-8). Yellow perch ranged in length from 3.5 to 10.1 inches, with an average length of 6.4 inches ($s = 2.0$ inches; $n = 39$), and an average weight of 0.1 pounds ($s = 0.1$ pounds; $n = 39$ Table 5-8).

Many fish collected in Lake Lory appeared in good condition. However, black spot, which is caused by a parasite (larval trematode) that burrows into the skin of the fish, was observed in several of the bluegills, black crappie, largemouth bass, smallmouth bass, and yellow perch.

Aquatic macroinvertebrate sampling was conducted on June 6, 2014 within Lake Lory where a total of 87 macroinvertebrates were collected (Table 5-9). True flies, dragonflies, and snails were the most abundant macroinvertebrates within Lake Lory. No threatened, endangered, or special concern macroinvertebrate species were observed in Lake Lory.

Floating, submerged, and emergent aquatic vegetation were observed in patches along the shoreline of Lake Lory. Burreed (*Sparganium* sp.), floating pondweed (*Potamogeton natans*), bigleaf pondweed (*Potamogeton amplifolius*), rushes, and water-starwort (*Callitriche* sp.) were the predominant species of aquatic vegetation that were observed along the shoreline (Photographs C-9 and C-10). Large woody debris in the form of

downed trees and submerged standing timber contributed to the aquatic habitat of Lake Lory.

Water temperature, pH, dissolved oxygen, and conductivity were measured on the surface of Lake Lory near the western shoreline on June 9, 2014. The surface water temperature (20.1 °C) and conductivity of Lake Lory was low (64 µS/cm; Table 5-6).

5.2.2 Wetland Complex EE

Wetland Complex EE was sampled for fish using a backpack shocker. A total of 17 fathead minnows (*Pimephales promelas*) were collected during the survey.

Aquatic macroinvertebrate sampling was conducted on June 06, 2014, where a total of 79 macroinvertebrates were collected (Table 5-10). Dragonflies, scuds, and snails were the most frequently collected macroinvertebrates within Wetland Complex EE. No threatened, endangered, or special concern macroinvertebrate species were observed in Wetland Complex EE.

Wetland Complex EE was predominantly vegetated with cattails and submerged patches of stonewort (*Chara* sp.; Photographs C-11 and C-12). The highest observed pH value (7.9) of this survey was recorded in the wetland (Table 5-6). Conductivity was the highest (378 µS/cm) relative to other sample stations (Table 5-6).

6.0 REFERENCES

- AEM, April 2007a. Humboldt Mill Aquatic Survey Report.
- AEM, October 2007b. Aquatic Survey of the Humboldt Tailings Disposal Facility.
- AEM, October 2007c. Humboldt Mill Project, Aquatic Survey of Wetland EE Complex.
- AEM, April 2008. Humboldt Mill Aquatic Survey Report 2007.
- AEM, March 2009. Humboldt Mill Aquatic Survey Report 2008.
- Bailey, R. M., W. C. Latta, and G. R. Smith. 2003. An atlas of Michigan fishes with keys and illustrations for their identification. Miscellaneous Publications, Museum of Zoology, No. 192, University of Michigan, Ann Arbor, MI.
- Becker, G. C. 1983. Fishes of Wisconsin. The University of Wisconsin Press, Madison, WI.
- Bright, E. 2014. "Aquatic Insects of Michigan." Museum of Zoology Insect Division and School of Natural Resources and Environment. University of Michigan, Ann Arbor, MI. Website, <<http://www.insects.ummz.lsa.umich.edu>> (Accessed 8/1/2014).
- Coon, T. G. 2001. Key to the fishes of Michigan. Michigan State University. East Lansing, MI.
- Cummings, K. S., and C. Mayer. 1992. Field guide to freshwater mussels of the Midwest. Illinois Natural History Survey, Champaign, IL.
- Merritt, R. W., K. W. Cummins, and M. B. Berg. 2008. An Introduction to the Aquatic insects of North America, 4th Edition. Kendall/Hunt Publishing Co., Dubuque, Iowa.
- Merritt, R. W., V. H. Resh, and K. W. Cummins. 1996. Design of aquatic insect studies: Collecting sampling and rearing procedures. *in* An Introduction to the Aquatic Insects of North America (2nd edition), Merritt, R. W., Cummins, K.W. editors. Kendall/Hunt Publishing Co., Dubuque, Iowa.
- MDEQ. 2008. GLEAS Procedure 51 - Qualitative biological and habitat survey protocols for wadable streams and rivers, April 24, 1990. Revised June 1991, August 1996, January 1997, May 2002, and December 2008. Michigan Department of Environmental Quality, Surface Water Quality Division, Lansing, MI.
- Michigan Natural Features Inventory. 2014. Marquette County Element Data. Michigan State University, East Lansing, MI.

Michigan Department of Natural Resources Fisheries Order DNR-DFI 101 FO-210.04. 2014.

Website, <http://www.michigandnr.com/law/law_book/orders/Fisheries%20Orders.html#FO210> (Accessed 9/15/2014).

Pennak, R. W. 1990. Freshwater invertebrates of the United States: protozoa to mollusca. 4th ed. John Wiley and Sons, Inc. 656 pp.


Schneider, J. C., G. R. Alexander, and J. W. Merna. 2000. Modules for lake and stream surveys. Chapter 2 *in* Schneider, J. C., editor, Manual of fisheries survey methods II: with periodic updates. Michigan Department of Natural Resources, Fisheries Special Report 25, Ann Arbor.

Smith-Root, Inc. 2004. Portable Electrofishers. Vancouver, WA.

Van Deventer, J. S., and W. S. Platts. 1983. Sampling and estimating fish populations from streams. Transactions of the North American Wildlife and Natural Resources Conference. 48: 349-354.

EXHIBIT A

REPORT FIGURES



Aerial imagery obtained from Michigan Center of Geographic Information (<http://www.michigan.gov/cgi/>)

Legend

- ! Humboldt Mill
- Roads




0 1.5 3 6 9 12 Miles

AeM

ADVANCED
ECOLOGICAL
MANAGEMENT

PROJECT	Humboldt Mill - Eagle Mine
TITLE	Humboldt Mill Location
FIGURE	1-1



Legend

- Wetland Complex EE Station
- Stream Sample Station Locations

0 875 1,750 3,500 5,250 7,000 Feet



AeM

ADVANCED
ECOLOGICAL
MANAGEMENT

PROJECT

Humboldt Mill - Eagle Mine


TITLE

Sample Station Locations

FIGURE

1-2

Aerial imagery obtained from Michigan Center for Geographic Information (<http://www.michigan.gov/cgi/>)



Aerial imagery obtained from Michigan Center of Geographic Information (<http://www.michigan.gov/cgi/>)

Legend

-] Fyke Net Locations
- # Gill Net Locations
- Electroshocker Transect Locations



PROJECT	Humboldt Mill - Eagle Mine
TITLE	Lake Lory Gear Locations
FIGURE	1-3

EXHIBIT B

REPORT TABLES

Table 1-1. Summary of the Procedure 51 Macroinvertebrate and Aquatic Habitat Scores for all Stream Stations, 2014.

Station Number	1	5	MBER1	MBER2
Fish Score Rating	Poor	Poor	Poor	Poor
Macroinvertebrate Score Rating	Acceptable	Acceptable	Acceptable	Acceptable
Stream Habitat Score Rating	Excellent	Good	Excellent	Excellent

Table 3-1. Stream Station Location Description.

Station Number	Stream Name	Latitude/Longitude NAD 1983	Township/Range/Section	Location Description
1	Unnamed Tributary of the Black River	N 46.47222 W 87.90249	Humboldt Twp. T47N, R29W, Sec 14	Upstream extent approximately 30 feet downstream of road crossing
5	Unnamed Tributary of the Middle Branch Escanaba River	N 46.49569 W 87.88276	Humboldt Twp. T47N, R29W, Sec 1	Extends between FX road and Highway 41
MBER1	Middle Branch Escanaba River	N 46.49899 W 87.88609	Humboldt Twp. T47N, R29W, Sec 1	Downstream extent approximately 2,683 feet upstream of Highway 41
MBER2	Middle Branch Escanaba River	N 46.527053 W 87.912157	Champion Twp. T48N, R29W, Sec 27	Upstream extent approximately 1,000 feet upstream of the former L.S. & I Railroad bridge

NAD 1983 = North American Datum of 1983

N = North

W = West

T = Township

R = Range

Sec = Section

Table 5-1. Stream Fish Collection Data – Stations 1, 5, MBER1 and MBER2.

Scientific Name	Common name	Station			
		1	5	MBER1	MBER2
<i>Catostomus commersonii</i>	White sucker			4	1
<i>Cottus bairdii</i>	Mottled sculpin			1	5
<i>Cualea inconstans</i>	Brook stickleback	1			
<i>Esox lucius</i>	Northern pike			7	2
<i>Lepomis macrochirus</i>	Bluegill			1	
<i>Margariscus margarita</i>	Pearl dace	8			
<i>Nocomis biguttatus</i>	Hornyhead chub				1
<i>Notropis cornutus</i>	Common shiner				4
<i>Percina flavescens</i>	Yellow perch			2	1
<i>Phoxinus eos</i>	Northern redbelly dace	13			
<i>Rhinichthys atratulus</i>	Blacknose dace			1	1
<i>Semolitus atromaculatus</i>	Creek chub				1
<i>Umbra limi</i>	Central mudminnow	9	1	1	3
Total Captured		31	1	17	19

Table 5-2. Stream Macroinvertebrate Collection Data – Stations 1, 5, MBER1 and MBER2.

TAXA	Station 1	Station 5	Station MBER1	Station MBER2
ANNELIDA (segmented worms)				
Hirudinea (leeches)	1	3	2	
Oligochaeta (worms)				
ARTHROPODA				
Crustacea				
Amphipoda (scuds)	35	11	10	3
Decapoda (crayfish)			11	5
Isopoda (sowbugs)	1	302	35	
Insecta				
Ephemeroptera (mayflies)				
Baetiscidae			2	
Baetidae		7		
Caenidae	1			
Ephemerellidae			1	5
Ephemeridae			1	
Heptageniidae			10	9
Siphonuridae			29	18
Odonata				
Anisoptera (dragonflies)				
Aeshnidae		2	2	1
Corduliidae	1	3		3
Gomphidae			2	
Libellulidae	1		1	1
Zygoptera (damselflies)				
Calopterygidae		5	4	3
Coenagrionidae	1			1
Plecoptera (stoneflies)				
Perlidae				1
Perlodidae			1	
Hemiptera (true bugs)				
Bellostomatidae			1	
Corixidae	2		1	3
Gerridae	2			
Megaloptera				
Sialidae (alder flies)			16	7
Trichoptera (caddisflies)				
Hydropsychidae		1		3
Lepidostomatidae				1
Limnephilidae	9	20	86	70
Molannidae			1	
Phryganeidae			1	1
Polycentropodidae			2	
Coleoptera (beetles)				
Dytiscidae (total)	1	6	9	2
Gyrinidae (adults)			3	

Table 5-2 (Continued). Stream Macroinvertebrate Collection Data – Stations 1, 5, MBER1 and MBER2.

TAXA	Station 1	Station 5	Station MBER1	Station MBER2
Haliplidae (adults)	1		1	1
Elmidae				1
Gyrinidae (larvae)		1		
Diptera (flies)				
Chironomidae	12	17	11	31
Simuliidae	3	4	6	226
Tipulidae			1	
MOLLUSCA				
Gastropoda (snails)				
Physidae		1		
Planorbidae	2	1		
Pelecypoda (bivalves)				
Pisidiidae	3			
Sphaeriidae		8		5
Total	76	392	250	401

Table 5-3. Stream Macroinvertebrate Scores and Community Ratings – Stations 1, 5, MBER1 and MBER2

METRIC	Station 1		Station 5		Station MBER1		Station MBER2	
	Value	Score	Value	Score	Value	Score	Value	Score
TOTAL NUMBER OF TAXA	16	0	15	1	27	0	23	0
NUMBER OF MAYFLY TAXA	1	-1	1	0	5	1	3	0
NUMBER OF CADDISFLY TAXA	1	-1	2	-1	4	0	4	0
NUMBER OF STONEFLY TAXA	0	-1	0	-1	1	0	1	0
PERCENT MAYFLY COMP.	1.32	0	1.79	0	17.20	0	7.98	0
PERCENT CADDISFLY COMP.	11.84	0	5.36	0	36.00	1	18.70	0
PERCENT CONTR. DOM. TAXON	46.05	-1	77.04	-1	34.40	-1	53.36	-1
PERCENT ISOPOD, SNAIL, LEECH	5.26	0	78.32	-1	14.80	-1	0.00	1
PERCENT SURF. AIR BREATHERS	7.85	0	1.53	1	6.00	0	1.50	1
TOTAL SCORE		-4		-2		0		1
COMMUNITY RATING		ACCEPTABLE		ACCEPTABLE		ACCEPTABLE		ACCEPTABLE

Table 5-4. Stream Habitat Scores and Ratings – Stations 1, 5, MBER1 and MBER2.

HABITAT METRIC	Station 1 glide/pool	Station 5 glide/pool	Station MBER1 glide/pool	Station MBER2 glide/pool
Substrate and Instream Cover				
Epifaunal Substrate/Avail. Cover	10	12	16	17
Pool Substrate Characterization	10	13	16	17
Pool Variability	11	5	16	15
Sediment Deposition	18	12	17	17
Channel Morphology				
Maintained Flow Volume	8	6	9	8
Flashiness	8	5	7	8
Channel Alteration	18	9	15	18
Channel Sinuosity	17	5	16	19
Riparian and Bank Structure				
Bank Stability (L)	9	9	10	9
Bank Stability (R)	9	9	8	9
Vegetative Protection (L)	10	9	10	10
Vegetative Protection (R)	10	9	8	10
Riparian Veg. Zone Width (L)	10	8	10	10
Riparian Veg. Zone Width (R)	10	9	8	9
Total Score	158	120	166	176
Habitat Rating	Excellent	Good	Excellent	Excellent

L = Left bank facing downstream**R = Right bank facing downstream**

Table 5-5. Stream Station Dimensions

Station	Length	Width (ft)		Depth (ft)	
		Average*	s	Average *	s
1	100	7.1 (3)	3.0	1.4 (9)	0.5
5	180	3.5 (3)	1.6	0.4 (9)	0.2
MBER1	1,000	36.9 (3)	2.6	3.2 (9)	0.7
MBER2	1,000	23.3 (3)	0.4	1.6 (9)	0.7

*sample size is indicated within ()

s = standard deviation

ft = feet

Table 5-6. Average Water Quality Parameters –Stations 1, 5, MBER1, MBER2 and Surface Water Bodies.

Station	Date	Time	Temperature °C	pH	Conductivity µS/cm	Percent Dissolved Oxygen	Dissolved Oxygen mg O ₂ /L	Discharge gpm
1	6/10/2014	8:25	13.6 (0.1)	6.9 (0.5)	154 (3.5)	34.4 (0.8)	5.0 (0.1)	861
5	6/09/2014	17:14	15.0 (0.1)	7.6 (0.3)	230 (8.5)	70.3 (4.9)	7.0 (0.5)	95
MBER1	6/06/2014	11:36	16.6 (0.1)	6.8 (0.5)	73 (2.6)	60.9 (0.9)	5.9 (0.1)	29,096
MBER2	6/06/2014	16:05	20.3 (0.6)	6.8 (0.2)	36 (0.0)	77.1 (3.8)	7.0 (0.3)	13,828
Lake Lory (Surface Water)	6/09/2014	13:55	20.1 (n.a.)	7.7 (n.a.)	64 (n.a.)	84.5 (n.a.)	7.7 (n.a.)	n.a.
Wetland Complex EE	6/09/2014	15:19	17.5 (n.a.)	7.9 (n.a.)	378 (n.a.)	99.6 (n.a.)	9.5 (n.a.)	n.a.

°C = degrees Celsius

µS/cm = microsiemens per centimeter

mg O₂/L = milligrams of oxygen per liter

gpm = gallons per minute

n.a. = not applicable – one sample point

standard deviation is indicated within ()

Table 5-7. Lake Lory Fish Collection Data.

Scientific Name	Common Name	Number of Taxa by Sample Gear			Total
		Fyke Nets	Gill Nets	Boom Shocker	
<i>Catostomus commersonii</i>	White sucker	2		3	5
<i>Etheostoma caeruleum</i>	Rainbow darter	1			1
<i>Hybognathus hankinsonii</i>	Brassy minnow	1			1
<i>Lepomis gibbosus</i>	Pumpkinseed sunfish	2		1	3
<i>Lepomis macrochirus</i>	Bluegill	29		40	69
<i>Micropterus dolomieu</i>	Smallmouth bass			15	15
<i>Micropterus salmoides</i>	Largemouth bass			79	79
<i>Notropis heterolepis</i>	Blacknose shiner	4			4
<i>Pimephales notatus</i>	Bluntnose minnow	3			3
<i>Perca flavescens</i>	Yellow perch		19	20	39
<i>Pomoxis nigromaculatus</i>	Black crappie	3	2	3	8
Total Collected by Gear		45	21	161	
Total Number of Fish Collected					227

Table 5-8. Lake Lory Fish Size.

Common Name	Length (inches)				Weight (pounds)			
	min	max	average*	s	min	max	average*	s
Black crappie	4.3	11.4	8.2(8)	2.7	0.04	0.72	0.5(5)	0.5
Blacknose shiner	2.2	2.5	2.4(4)	0.1	0.003	0.004	0.00(4)	0.003
Bluegill	0.4	9.3	5.5(69)	2.0	0.01	0.49	0.14(69)	0.1
Bluntnose minnow	2.2	3.1	2.6(3)	0.5	0.004	0.01	0.01(7)	0.003
Brassy minnow	2.9	2.9	n.a.(1)	n.a.	0.01	0.01	n.a.(1)	n.a.
Largemouth bass	5.9	17.3	12.0(79)	1.9	0.08	2.76	0.82(79)	0.4
Pumpkinseed sunfish	4.0	6.6	5.0(3)	1.5	0.05	0.29	0.13(3)	0.1
Rainbow darter	1.8	1.8	n.a.(1)	n.a.	0.002	0.002	n.a.(1)	n.a.
Smallmouth bass	8.3	14.6	11.5(15)	2.2	0.25	1.25	0.7(15)	0.4
White sucker	6.1	14.6	10.0(5)	3.6	0.08	1.22	0.5(5)	0.5
Yellow perch	3.5	10.1	6.4(39)	2.0	0.01	0.37	0.12(39)	0.1

* sample size is indicated within ()

s = standard deviation

min = minimum

max = maximum

n.a. = not applicable

Table 5-9. Lake Lory Macroinvertebrates.

Order	Family	Genus	Taxa Count
Amphipoda	Hyalellidae	<i>Hyalella</i>	6
Basommatophora	Lymnaeidae	<i>Stagnicola</i>	1
Basommatophora	Planorbidae	<i>Helisoma</i>	6
Coleoptera	Elmidae		1
Coleoptera	Gyrinidae	<i>Gyrinus</i>	2
Coleoptera	Halplidae	<i>Peltodytes</i>	3
Decapoda	Cambaridae		2
Diptera	Chaoboridae	<i>Chaoborus</i>	2
Diptera	Chironomidae		21
Ephemeroptera	Caenidae	<i>Caenis</i>	2
Ephemeroptera	Ephemerellidae	<i>Attenella</i>	6
Mesogastropoda	Hydrobiidae	<i>Marstonia</i>	17
Odonata	Coenagrionidae	<i>Coenagrion</i> , <i>Enallagma</i>	11
Odonata	Corduliidae	<i>Epithea</i>	1
Odonata	Corduliidae	<i>Somatochlora</i>	1
Odonata	Gomphidae	<i>Hagenius</i>	1
Rhynchobdellida	Glossiphoniidae	<i>Helobdella</i>	1
Trombidiformes	Hydrachnidae	<i>Hydracarina</i>	2
Veneroida	Pisidiidae	<i>Pisidium</i>	1
Total			87

Table 5-10. Wetland Complex EE Macroinvertebrates.

Order	Family	Genus	Taxa Count
Amphipoda	Hyalellidae	<i>Hyalella</i>	18
Basommatophora	Lymnaeidae	<i>Stagnicola</i>	3
Basommatophora	Physidae	<i>Physa</i>	4
Basommatophora	Planorbidae	<i>Helisoma</i>	12
Coleoptera	Dytiscidae	<i>Coptotomus</i>	1
Coleoptera	Dytiscidae	<i>Derovatellus</i>	1
Coleoptera	Gyrinidae	<i>Gyrinus</i>	1
Coleoptera	Hydrophilidae	<i>Tropisternus</i>	1
Diptera	Ceratopogonidae	<i>Bezzia/Palpomyia</i>	1
Diptera	Chironomidae		4
Diptera	Culicidae		1
Ephemeroptera	Baetidae	<i>Callibaetis</i>	1
Ephemeroptera	Caenidae	<i>Caenis</i>	5
Hemiptera	Macroveliidae		1
Odonata	Aeshnidae	<i>Anax</i>	1
Odonata	Coenagrionidae	<i>Coenagrion, Enallagma</i>	13
Odonata	Libellulidae	<i>Leucorrhina</i>	2
Odonata	Libellulidae	<i>Libellula</i>	3
Trichoptera	Limnephilidae	<i>Limnephilus</i>	2
Veneroida	Pisidiidae	<i>Pisidium</i>	4
Total			79

EXHIBIT C

PHOTOGRAPHS



Photograph C-1. Station 1 – Upstream Extent. View looking south, downstream.



Photograph C-2. Station 1 – Downstream Extent. View looking west, upstream.



Photograph C-3. Station 5 – Upstream Extent. View to north, downstream.



Photograph C-4. Station 5 – Downstream Extent. View to south, upstream.



Photograph C-5. Station MBER1– Upstream Extent. View to east, downstream.



Photograph C-6. Station MBER1 – Downstream Extent. View to west, upstream.



Photograph C-7. Station MBER2 – Upstream Extent. View to south, downstream.



Photograph C-8. Station MBER2 – Downstream Extent. View to north, upstream.



Photograph C-9. Lake Lory. View to north.



Photograph C-10. Lake Lory. View to south.



Photograph C-11. Wetland Complex EE North of HTDF. View to northwest.



Photograph C-12. Wetland Complex EE North of HTDF. View to north.